



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 300 466 B1

(12)

EUROPEAN PATENT SPECIFICATION

- (43) Date of publication of patent specification: 13.09.95 (51) Int. Cl.®: C12N 15/09, C12N 1/15, C12N 15/00
- (21) Application number: 88111713.9
- (22) Date of filing: 20.07.88

The file contains technical information submitted after the application was filed and not included in this specification

- (54) Characterization and structure of genes for protease A and protease B from streptomyces griseus.

- (26) Priority: 21.07.87 CA 542678
- (43) Date of publication of application: 25.01.89 Bulletin 89/04
- (43) Publication of the grant of the patent: 13.09.95 Bulletin 95/37
- (24) Designated Contracting States: AT BE CH DE ES FR GB GR IT LI LU NL SE
- (56) References cited:
EP-A- 0 161 629 EP-A- 0 179 449
EP-A- 0 187 630 EP-A- 0 196 375
EP-A- 0 213 898 EP-A- 0 222 279

JOURNAL OF BACTERIOLOGY, vol. 169, no. 8, August 1987, American Society for Microbiology; G. HENDERSON et al., pp. 3778-3784

- (23) Proprietor: Cangene Corporation
6280 Northwest Drive
Mississauga
Ontario L4V 1J7 (CA)
- (72) Inventor: Garvin, Robert T.
33 Deforest Road

Toronto, ON, M6S 1J1 (CA)
Inventor: Henderson, Graham
76 Doncaster Drive
Bramalea, ON, L6T 1T1 (CA)
Inventor: Krygsman, Phyllis
113 Woodview Avenue
Pickering, ON, L1V 1L1 (CA)
Inventor: Liu, Ci Jun
695 Martin Grove Road,
No. 512
Weston, ON, M9R 3L6 (CA)
Inventor: Davey, Cheryl
175 Dinnick Crescent
Toronto, ON, M4N 1M4 (CA)
Inventor: Malek, Lawrence T.
4 Sproule Drive
Brampton, ON, L6V 4A5 (CA)

- (24) Representative: Perry, Robert Edward et al
GILL JENNINGS & EVERY
Broadgate House
7 Eldon Street
London EC2M 7LH (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

FIELD OF THE INVENTION

This invention relates to a biologically pure DNA signal sequence which encodes an amino acid signal peptide necessary for directing the secretion from certain defined hosts of proteins in bioactive form.

BACKGROUND OF THE INVENTION

In the biological production of commercially viable proteins by the fermentation of microorganisms, the ability to produce the desired proteins by fermentation with secretion of the proteins by the microorganisms into the broth is very significant. However, there are many commercially viable proteins encoded by genetically engineered DNA constructs which are not secreted by the cells in which the DNA is expressed. This often necessitates harvesting the cells, bursting the cell walls, recovering the desired proteins in pure form and then chemically re-naturing the pure material to restore its bioactive function. This downstream processing, as it is called, is illustrated in Figure 1.

Some cells and microorganisms carry out the biological equivalent of downstream processing by secreting proteins in bioactive form. The mechanism which directs the secretion of some proteins through the cell walls is not fully understood. For example, in *Streptomyces griseus*, an organism used for the commercial production of Pronase, the species secretes many extra cellular proteins (Jurasek, L., P. Johnson, R.W. Olafson, and L.B. Smillie (1971), An improved fractionation system for pronase on CM-Sephadex *Can. J. Biochem.*, 49:1195-1201). Protease A and protease B, two of the serine proteases secreted by *S. griseus*, have sequences which are 61% homologous on the basis of amino acid identity (Fujinaga, M., L.T.J. Delbaere, G.D. Brayer, and M.N.G. James (1985), Refined structure of α -lytic protease at 1.7 Å resolution; Analysis of hydrogen bonding and solvent structure, *J. Mol. Biol.*, 183:479-502; Jurasek, L., M.R. Carpenter, L.B. Smillie, A. Gertler, S. Levy, and L.H. Ericsson (1974), Amino acid sequencing of *Streptomyces griseus* protease B, A major component of pronase, *Biochem. Biophys. Res. Comm.*, 61:1095-1100; Young, C.L., W.C. Barker, G.M. Tomaseff, and M.O. Dayhoff (1978), Serine proteases, in M.O. Dayhoff (ed.), *Atlas of Protein Sequence and Structure* 5, suppl. 3:73-83). These proteases also have similar tertiary structure, as determined by X-ray crystallography (Delbaere, L.T.J., W.L.B. Hutcheon, M.N.G. James, and W.E. Thiessen (1975), Tertiary structural differences between microbial serine proteases and pancreatic serine enzymes, *Nature* 257:758-763; Fujinaga, M., L.T.J. Delbaere, G.D. Brayer, and M.N.G. James (1985), Refined structure of α -lytic protease at 1.7 Å resolution; Analysis of hydrogen bonding and solvent structure, *J. Mol. Biol.*, 183:479-502; James, M.N.G., A.R. Sielecki, G.D. Brayer, L.T.J. Delbaere, and C.-A. Bauer (1980), Structures of product and inhibitor complexes of *Streptomyces griseus* protease A at 1.8 Å resolution, *J. Mol. Biol.*, 144:43-85). Although the structures of proteases A and B have been extensively studied, the genes encoding these proteases have not been characterized before. EP-A-0 222 279 discloses signal peptides derived from *Streptomyces*.

SUMMARY OF THE INVENTION

In accordance with this invention, the genes encoding protease A and protease B of *S. griseus* have been isolated and investigated to reveal DNA sequences which each direct the secretion of an encoded protein fused either directly or indirectly to a signal peptide encoded by the DNA.

According to an aspect of the invention, a recombinant DNA sequence comprises a signal sequence and a gene sequence encoding a protein. The recombinant DNA sequence, when expressed in a living cell, encodes an amino acid signal peptide with the protein. The signal peptide directs secretion of the protein from a cell within which the DNA signal sequence is expressed.

According to another aspect of the invention, a biologically pure isolated DNA signal sequence encodes a 38 amino acid signal peptide which directs secretion of a recombinant gene-sourced protein linked to such 38 amino acid signal peptide, from a cell in which the DNA signal sequence is expressed. The DNA signal sequence is isolated from *Streptomyces griseus*.

According to another aspect of the invention, the DNA signal sequence in conjunction with a gene sequence encoding a protein is inserted into a vector, such as a plasmid or a phage.

According to another aspect of the invention, the DNA signal sequence is adapted for expression in a living cell having enzymes catalyzing the formation of disulphide bonds.

According to another aspect of the invention, the biologically pure isolated DNA signal sequence of Figure 4a.

According to another aspect of the invention, the biologically pure isolated DNA signal sequence of Figure 5a.

According to another aspect of the invention, a fused protein is encoded by the recombinant DNA sequence of Figure 4 or Figure 5.

5 According to another aspect of the invention, a transformed prokaryotic cell is provided which has inserted therein a suitable vector including the recombinant DNA encoding the signal protein. The transformed prokaryotic cell may be selected from the *Streptomyces* genera.

According to another aspect of the invention, a biologically pure culture has a transformed prokaryotic cell with the recombinant DNA sequence in a suitable vector. The culture is capable of producing, as an intermediate, the fused protein of the amino acid signal peptide and the protein. The protein itself is produced in a recoverable quantity upon fermentation of the transformed cell in an aqueous nutrient medium. The signal peptide directs secretion of the protein from the cell.

According to another aspect of the invention, a biologically pure culture, transformed with the functional signal sequence as described above, is able to direct the secretion from the cell of proteins whose bioactivity is dependent upon the formation of correctly positioned intramolecular disulphide bonds.

75 A biologically pure DNA sequence encoding a fused protein including protease A has the combined DNA sequence of Figures 4a, 4b and 4c.

A biologically pure DNA sequence encoding a fused protein including protease B has the combined DNA sequence of Figures 5a, 5b and 5c.

20

BRIEF DESCRIPTION OF THE DRAWINGS

With reference to the Figures, a variety of short forms have been used to identify restriction endonucleases, amino acids, deoxyribonucleic acids and related information. Standard nomenclature has been used in identifying all of these components as are readily appreciated by those skilled in the art.

Preferred embodiments of the invention are described with respect to the drawings, wherein:

Figure 1 illustrates downstream processing;

Figure 2 shows restriction endonuclease maps of DNA fragments of *sprA* and *sprB*;

Figure 3 illustrates restriction endonuclease maps and sequencing strategies in sequencing DNA fragments containing *sprA* and *sprB*;

30 Figure 4 is the DNA sequence of *sprA*;

Figure 4a is the DNA sequence encoding the *sprA* (protease A) signal peptide;

Figure 4b is the DNA sequence encoding the *sprA* (protease A) propeptide;

Figure 4c is the DNA sequence encoding mature protease A;

35 Figure 5 is the DNA sequence of *sprB*;

Figure 5a is the DNA sequence encoding the *sprB* (protease B) signal peptide;

Figure 5b is the DNA sequence encoding the *sprB* (protease B) propeptide;

Figure 5c is the DNA sequence encoding mature protease B;

40 Figure 6 is an alignment of the amino acid sequences deduced from *sprA* and *sprB* to develop homology between the two sequences.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The organism *Streptomyces griseus* is a well recognized microorganism. It is commercially used for the production of the enzyme Pronase. It is appreciated, however, that this organism also secretes two enzymes, protease A and protease B, which are both serine proteases. Although the structure of proteases A and B have been extensively studied, the genes encoding these proteins, and the manner in which this genetic information is used to signal secretion by the cells, is not understood. According to this invention, the genes which encode protease A and protease B and provide for the secretion of these proteins in bioactive form have been discovered. It has been determined that each of protease A and B is included in a precursor protein which is processed to remove an amino-terminal polypeptide portion from the mature protease. It has further been determined that each of protease A and B precursor proteins is enzymatically processed to form correctly-positioned intramolecular disulphide bonds, which processing is concomitant with removal of the amino terminal addressing peptide from the mature precursor. The discovered genes, which encode proteases A and B, their intermediate address-competent forms, and their control elements, have been designated *sprA* and *sprB*.

As discussed in the following articles, Jurasek, L., M.R. Carpenter, L.B. Smillie, A. Gertler, S. Levy, and L.H. Ericsson (1974), Amino acid sequencing of *Streptomyces griseus* protease B, a major component of

protease, Biochem. Biophys. Res. Comm. 61:1095-1100; Young, C.L., W.C. Barker, C.M. Tomaselli, and M.O. Dayhoff (1978), Serine proteases, In M.O. Dayhoff (ed.), Atlas of Protein Sequence and Structure 5, suppl. 3:73-83, proteases A and B are homologous proteins containing several segments of identical amino acid sequence. In accordance with this invention, the genetic code, which makes and directs the secretion of each of proteases A and B, has identical DNA sequences corresponding to the regions of identity for the homologous proteins proteases A and B. In order to isolate the genes, this assumption, that identity in portions of the gene sequences would occur, was made so that an oligonucleotide probe could be designed from one of the similar regions in the sequences.

In order to extrapolate the gene sequence which would encode the similar amino acid sequence, the known codon bias for *Streptomyces* was relied upon to develop the nucleotide probe (see Berman, V., D. Filipula, W. Harber, M. Bibb, and E. Katz (1985), The nucleotide sequence of the tyrosinase gene from *Streptomyces antibioticus* and characterization of the gene product, Gene 37:101-110; Bibb, M.J., M.J. Bibb, J.M. Ward, S.N. Cohen (1985), Nucleotide sequences encoding and promoting expression of three antibiotic resistance genes indigenous to *Streptomyces*, Mol. Gen. Genet. 199:26-38; (Thompson, C.J., and G.S. Gray (1983), Nucleotide sequence of a streptomycete aminoglycoside phosphotransferase gene and its relationship to phosphotransferases encoded by resistance plasmids, Proc. Natl. Acad. Sci. USA 80:5190-5194). Once the probe was constructed, it was then possible to probe the DNA sequences of *S. griseus* to determine if there were any corresponding nucleic acid sequences in the microorganism. Since it was known that there were two proteases, A and B, the oligonucleotide probe should have revealed two DNA fragments detected by hybridization analysis, and in fact, not only did the probe hybridize equally to two fragments generated in the genomic library of *S. griseus*, but also two fragments generated by *Bam*HI digest (8.4 kb and 6.8 kb) or *Eco*III (11 kb and 2.8 kb) were isolated from the genomic library. As a cross-check with respect to the predictability of such probe, the same fragments were detected in genomic DNA libraries of other isolates of *S. griseus*. It was noted, however, that there was no such hybridization of the oligonucleotide probe with DNA from other *Streptomyces* such as *S. lividans*.

Plasmids were constructed containing digested fragments of *S. griseus*. The oligonucleotide probe was used to isolate developed plasmids containing *sprA* and *sprB*. The screening by use of the probe was accomplished by colony blot hybridization where approximately 15,000 *E. coli* transformants containing the developed plasmids were screened. Twelve transformants were detected by the probe and isolated for further characterization. These colonies contained two distinct classes of plasmid based on restriction analysis. As determined from the hybridization of genomic DNA, the plasmids contained either the 6.8 kb or the 8.4 kb *Bam*HI fragment. These fragments contained the *sprA* and *sprB* genes.

The fragments as isolated by hybridization screening were tested for the expression of proteolytic activity. With these plasmids identified, such characterization may be accomplished in accordance with a variety of known techniques in accordance with a preferred embodiment of this invention.

The 6.8 kb and 8.4 kb *Bam*HI fragments were ligated into the *Eco*III site of the vector pIJ702. Transformants of *S. lividans* containing these constructions were tested on a milk plate for secretion of proteases. A clear zone, which represented the degradation of the milk proteins, surrounded each transformant that contained either *Bam*HI fragment. It was noted that the clear zones were not found around *S. lividans* colonies which contained either pIJ702 only or no plasmid construct.

Proteolytic activity was also observed when the *Bam*HI fragments were cloned in either orientation with respect to the vector, thereby minimizing the possibility of read-through transcription of an incomplete protease gene. This observation provides evidence that the two *Bam*HI fragments contain an intact protease gene which is capable of effecting secretion in a different *Streptomyces* species, as for example the *S. lividans*. With this particularly relevant characterization of the *Bam*HI fragment, and knowing that the desired gene was in these fragments, it was possible to isolate and to sequence the genes encoding protease A and protease B.

According to a preferred aspect of this invention, the particular protease gene contained within each cloned *Bam*HI fragment was determined by dideoxy sequencing of the plasmids using the oligonucleotide probe as a primer in such analysis. The 8.4 kb *Bam*HI fragment was found to contain *sprB*, because a polypeptide deduced from the DNA sequence matched a unique segment of the known amino acid sequence of protease B. The 6.8 kb *Bam*HI fragment contained the *sprA* by process of elimination. The protease genes in these fragments were localized by digesting the plasmids and determining which of the restriction fragments of the plasmids were capable of hybridizing to the oligonucleotide probe.

Figure 2 shows detailed restriction maps of the 6.8 kb and 8.4 kb *Bam*HI fragments. Hybridization to the oligonucleotide probe was confined to a 0.9 kb *Pvu*II-*Stu*I fragment of *sprA* and a 0.6 kb *Pvu*II-*Pvu*I fragment of *sprB*. Such hybridization is indicated by the heavy lines in Figure 2. Hybridization to the cloned *Bam*HI fragments and the 2.8 kb *Eco*III fragment of *sprB* agrees with the hybridization to *Bam*HI and *Eco*III fragments

of genomic DNA. Thus, rearrangement of the BamHI fragments containing the protease genes is unlikely to have occurred.

The functional portions of the sprA- and sprB-containing DNA were determined by subcloning restriction fragments thereof into pJ702. The constructed plasmids were transformed into S. lividans and tested for proteolytic activity. The 3.2 kb BamHI- BglII fragment of sprA and the 2.8 kb BglII fragment of sprB, when subcloned into pJ702 in either orientation, resulted in the secretion of protease from S. lividans. The intact protease genes were further delimited to a 1.9 kb StuI fragment for sprA and a 1.4 kb BssHI fragment for sprB. With reference to Figure 2, each of these functionally active subclones are indicated below the restriction maps which contain the region for each gene which hybridized to the oligonucleotide probe.

In order to determine the nucleic acid sequence of the protease genes, the 3.2 kb BamHI-BglII fragment of sprA and the 2.8 kb BglII fragment of sprB were subcloned into pUC18 to facilitate further structural characterization. As shown in Figure 3, the restriction maps of these subclones and the strategies which were used to sequence the 1.4 kb SalI fragment containing sprA and the 1.4 kb BssHI fragment containing sprB are shown. The resultant DNA sequences of sprA and sprB are shown in Figures 4 and 5, respectively. The predicted amino acid sequence of protease A differed from the published sequence by the amidation of amino acid 133, whereas that of protease B was identical to the published sequence, (see Fujinaga, M., L.T.J. Delbaere, G.D. Brayer, and M.N.G. James (1985). Refined structure of α -lytic protease at 1.7 Å resolution; Analysis of hydrogen bonding and solvent structure. J. Mol. Biol. 183:479-502).

Analyzing the sequences of Figures 4 and 5, it is apparent that each sequence contains a large open reading frame with the coding region of the mature protease situated at the 3' end. For the protease A and protease B genes, the sequence encoding the carboxy-terminus of the protease is followed immediately by a translation stop codon. At the other end of the sequence, the predicted amino acid sequences appear to extend beyond the amino-termini of the mature proteases A and B by an additional 116 amino acids for sprA of Figure 4 and 114 amino acids for sprB of Figure 5. The putative GTG initiation codons at each of these positions (-116 for Figure 4; -114 for Figure 5) are each preceded by a potential ribosome binding site (as indicated by the series of five dots above the sequence) and followed by a sequence which encodes a signal peptide. The processing site for the signal peptidase (identified by the light arrow in Figures 4 and 5) is predicted at 38 amino acids from the amino-terminus of the putative precursor. [For clarity, that part of the nucleic acid sequences of Figures 4 and 5 corresponding to the signal peptide portion of sprA and sprB is reproduced in Figures 4A and 5A, respectively]. The propeptide is encoded by the remaining sequence between the signal processing site (light arrow) and the start of the mature protein (indicated at the dark arrow). [For clarity, that part of nucleic acid sequences of Figures 4 and 5 corresponding to the propeptide portion of sprA and sprB is reproduced in Figures 4B and 5B, respectively]. The mature protease is encoded by the codon sequence 1 through 181 for Figure 4 and 1 through 185 for Figure 5. [For clarity, that part of the nucleic acid sequences of Figures 4 and 5 corresponding to the mature protein portion of sprA and sprB is reproduced in Figures 4C and 5C, respectively]. The amino acid sequence for codons -116 through +181 of Figure 4 and the amino acid sequence for codons -114 through +185 of Figure 5, when made in the living cell S. griseus, are acted upon in a manner to produce in the culture medium externally of the living cells the mature bioactive enzymes protease A and protease B. The processing involved in accordance with the contained information encoded by that portion of the gene from start of the promoter to start of the mature protein in each case included providing a secretory address, the correct signal peptide processing site, the necessary propeptide structure not only for secretion but also for correct disulphide bond formation concomitant with secretion, and competent secretion in bioactive form.

In accordance with this invention, the ability of the signal peptide to direct the secretion of bioactive protein was established by inserting known DNA sequences at the beginning and at the end of known sequences. For example, consider the sequence shown in Figure 5. In particular, the promoter and initiator ATG of the aminoglycoside phosphotransferase gene, (Thompson, C.J., and G.S. Gray (1983). Nucleotide sequence of a streptomycete aminoglycoside phosphotransferase gene and its relationship to phosphotransferases encoded by resistance plasmids, Proc. Natl. Acad. Sci. USA, 80:5190-5194) had been inserted preceding the second codon (AGG at -113) of the signal sequence of Figure 5. Due to the insertion of this new promoter and initiator, the sprB gene, now under the control of this non-native promoter, directed both elevated levels and earlier expression of proteolytic activity when compared with the unaltered sprB gene. The secretion of bioactive protease B in this construction indicated that nucleic acid sequences preceding the GTG initiation codon at -114 are not required for the correct secretion of the protease B in bioactive form, provided an active and competent promoter is placed in the precise location indicated.

In order further to demonstrate the universality of the discovered signal peptide, the sprB coding region was replaced with a gene sequence encoding the mature amylase from S. griseus. Hence the nucleic acid sequence encoding the amylase was inserted in place of the sequence of Figure 5 to the right of the light

arrow. It was determined that the resulting genetic construction directed the production of an extracellular protein having an N-terminal alanine, properly positioned intramolecular disulphide bonds, and exhibiting amylolytic activity at a level comparable to that of a similar construction with the natural signal peptide of amylase. In accordance with this invention, the 38 amino acid signal peptide of Figures 4 and 4A and 5 and 5A is sufficient to direct the secretion of non-native protein in bioactive form.

Since both signal sequences encode for the signal peptides of Figures 4 and 4A and 5 and 5A, the organization of the coding regions of *sprA* and *sprB* were investigated by comparing the amino acid homology of the encoded peptide sequences. Such comparisons are set out in Figure 6 where amino acid homology has been compared for the signal peptide of Figure 6a, the propeptide of Figure 6b and the mature protease of Figure 6c. A summary of such homology is provided in the following Table I.

TABLE I

Homology of <i>sprA</i> and <i>sprB</i> Coding Regions			
	Length (codons)	Protein Homology %	DNA Homology %
Signal	38	50	58
Propeptide	79	43	62
NT protease ^a	87	45	58
CT protease ^b	103	75	75
Total Protease	190	61	67
Total coding region	307	55	65

a amino-termini of mature proteases (amino acids 1-87)

b carboxy-termini of mature proteases (amino acids 68-190)

The alignment of amino acid sequences translated from the coding regions of the *sprA* and *sprB* genes indicates an overall homology of 54% on the basis of amino acid identity. As indicated in Table I, the sequence homology is not uniformly distributed throughout the coding region of the *sprA* and *sprB* genes. The carboxy-terminal domains of the proteases A and B are 75% homologous as noted under the heading "CT protease" whereas the average homology for the remainder of the coding region is only 45%, indicated under the heading "NT protease". The amino terminal domains containing the signal and propeptide regions were similar in both extent of homology and distribution of consensus sequences, as indicated under the headings "signal" and "propeptide". The unexpectedly high DNA sequence homology relative to that of the protein sequences is particularly due to the 61% conservation in the third position of each codon of the sequence. These investigations, revealing the close homology between *sprA* and *sprB* genes, suggest that both genes originated by duplication of a common ancestral gene. With appropriate care and investigation, the commonality of the signal peptides can be determined, thus establishing the cue for secretion of proteins and hence providing sufficient information to construct, from the signal DNA of *sprA* and *sprB*, a single nucleic acid sequence which will be competent to direct protein secretion.

In accordance with the invention, a recombinant DNA sequence can be developed which encodes for desired protein where the expressed protein, in conjunction with the signal peptide and optionally the propeptide, provide for secretion of the desired protein in bioactive form. The recombinant DNA sequence may be inserted in a suitable vector for transforming a desired cell for manufacturing the protein. Suitable expression vectors may include plasmids and viral phages. As is appreciated by those skilled in the art, the bioactivity of secretory proteins is assured by establishing the correct configuration of intramolecular disulphide bonds. Thus, suitable prokaryotic hosts may be selected for their ability to display enzymatic activity of a type typified by, but not limited to, that of protein disulphide oxidoreductase, EC 5.3.4.1.

The particular protein encoded by the recombinant DNA sequence may include eukaryotic secretory enzymes, such as prochymosin, chymotrypsin, trypsin, amylases, ligninases, chymosin, elastases, lipases, and cellulases; prokaryotic secretory enzymes such as glucosyl isomerase, amylases, lipases, pectinases, cellulases, proteinases, oxidases, ligninases; blood factors, such as Factor VIII and Factor IX and Factor VIII-related biosynthetic blood coagulant proteins; tissue-type plasminogen activator; hormones, such as proinsulin; lymphokines, such as beta and gamma-interferon, and interleukin-2; enzyme inhibitors, such as extracellular proteins whose action is to destroy antibiotics either enzymatically or by binding, for example, a β -lactamase inhibitor, a trypsin inhibitor; growth factors, such as organism or nerve growth factors, epidermal growth factors, tumor necrosis factors, colony stimulating factors; immunoglobulin-related molecules, such as synthetic, designed, or engineered antibody molecules; cell receptors, such as cholesterol

receptor; viral molecules, such as viral hemagglutinins, AIDS antigen and immunogen, hepatitis B antigen and immunogen, foot-and-mouth disease virus antigen and immunogen; bacterial surface effectors, such as protein A; toxins such as protein insecticides, algicides, fungicides, and biocides; and systemic proteins of medical importance, such as myocardial infarct protein (MIP), weight control factor (WCF), caloric rate protein (CRP) and hirudin (HRD).

One skilled in the art can easily determine whether the use of any known or unknown organism will be within the scope of this invention in accordance with the above discussion and the following examples.

Microorganisms which may be useful in this respect as potential prokaryotic expression hosts include:

Order:

- 10 Actinomycetales; Family: Actinomycetaceae Genus: Matricornema, Lactiophora; Family Actinobacteria:
Genus Actinomyces, Agromyces, Arachina, Arcanobacterium, Arthrobacter, Brevibacterium, Cellulomonas,
Curtobacterium, Microbacterium, Oerskovia, Promicromonospora, Renibacterium, Rothia; Family Ac-
tinoplanetes; Genus Actinoplanes, Dactylosporangium, Micromonospora; Family Nocardiform ac-
tinomyces; Genus Caseobacter, Corynebacterium, Mycobacterium, Nocardia, Rhodococcus; Family Strep-
15 tomycetes; Genus Streptomyces, Streptovericillium; Family Maduromycetes; Genus Actinomadura, Excel-
lospora, Microspora, Planospora, Spirillospora, Streptosporangium; Family Thermospora; Genus Actinosyn-
nema, Nocardopsis, Thermophilla; Family Microspora; Genus Actinospora, Saccharospora; Family Ther-
moactinomycetes; Genus Thermoactinomyces; and the other prokaryotic genera: Acetivibrio, Acetobacter,
Achromobacter, Acinetobacter, Aeromonas, Bacterionema, Bilidobacterium, Flavobacterium, Kurthia, Lac-
20 tobacillus, Leuconostoc, Mycobacteria, Propionibacterium.

The following species from the genus Streptomyces are identified as particularly suitable as hosts:

- acidophilus, albus, amylophilus, argentiolus, aureofaciens, aureus, candidus, cellosolus, cellulolyticus,
coelicolor, creamorus, diastolicus, farinosus, flaventus, flavogriseus, fradiae, fulvoviridis, fungicidicus,
gelaticus, glaucescens, globisporus, griseolus, griseus, hygroscopicus, ligninolyticus, lipolyticus, lividans,
25 moderatus, olivochromogenus, parvus, phaenochromogenes, piliculus, proteolyticus, reclus, roseolus,
roseoviolaceus, scabies, thermofyticus, tumorstalicus, venezuelae, violaceus, violaceus-ruber, violaceus,
and viridochromogenes.
acrimycin
albioniger
30 ambifaciens
antibioticus
aspergilloides
charfreusis
clavuligerus
35 diastatochromogenes
echinatus
erythraeus
fendae
griseofuscus
40 kanamyceticus
kasugaensis
koganeiensis
lavendulae
parvulus
45 peuceilus
reticuli
rimosus
vinaceus

Also, the following eukaryotic hosts are potentially useful in the practice of this invention:

- 50 Absidia, Acremonium, Acrophialopora, Acrospeira, Altermaria, Arthrotrix, Ascotricha, Aureobasidium,
Beauveria, Bispora, Bjerkandera, Calocera, Candida, Cephalophora, Cephalosporium, Cerinomyces,
Chaetomium, Chrysosporium, Circinella, Cladosporium, Clonostix, Coccospora, Cochliobolus, Cunninghamella,
Curvularia, Cuslingophora, Dacrymyces, Dacryopinax, Dendryphon, Dictosporium, Doratomyces,
Drechslera, Eupenicillium, Flammulina, Fusarium, Gliocladium, Gliomastix, Graphium, Hansenula,
55 Humicola, Hyalodendron, Isaria, Klueckera, Kluyveromyces, Lipomyces, Mammaria, Merulius, Microascus,
Monodictys, Monosporium, Morchella, Mortierella, Mucor, Myceliophthora, Myrothecium, Neurospora,
Oedocephalum, Oidiendron, Pachysolen, Papularia, Papulaspora, Penicillium, Peniophora, Periconia,
Phaeoconiolellus, Phanerochaete, Phialophora, Piptocephalis, Pleurotus, Preussia, Pycnosporus, Rhion-

cladiella, Rhizomucor, Rhizopus, Rhodotorula, Robillarda, Saccharomyces, Schwanniomyces, Scolecobasidium, Scopulariopsis, Scytalidium, Stachybotrys, Tetraculium, Thamnidium, Thermioascus, Thermomyces, Thielavia, Tolypocladium, Torula, Torulopsis, Trametes, Tricellula, Trichocladium, Trichoderma, Trichurus, Truncatella, Ulocladium, Ustilago, Vericillium, Wardomyces, Xylogone, Yarrowia.

Preferred embodiments of the invention are exemplified in the following procedures. Such procedures and results are by way of example and are not intended to be in any way limited to the scope of the claims.

PREPARATIONS

Strains and Plasmids

Streptomyces griseus (ATCC 15395) was obtained from the American Type Culture Collection. Streptomyces lividans 68 (Bibb, M.J., J.L. Scholte, and S.N. Cohen (1980), A DNA cloning system for interspecies gene transfer in antibiotic-producing Streptomyces, Nature 284:528-531) and the plasmids pJ81 and pJ702 from the John Innes Institute; Thompson, C.J., T. Kieser, J.M. Ward, and D.A. Hopwood (1982), Physical analysis of antibiotic-resistance genes from Streptomyces and their use in vector construction, Gene 20:51-62; Katz, E., C.J. Thompson, and D.A. Hopwood (1983), Cloning and expression of the tyrosinase gene from Streptomyces antibioticus in Streptomyces lividans, J. Gen. Microbiol., 129:2703-2714). E. coli strain HB101 (ATCC 33694) was used for all transformations. Plasmids pUC8, pUC18 and pUC19 were purchased from Bethesda Research Laboratories.

Media, Growth and Transformation

Growth of Streptomyces mycelium for the isolation of DNA or the preparation of protoplasts was as described in Hopwood, D.A., M.J. Bibb, K.F. Chater, T. Kieser, C.J. Bruton, H.M. Kieser, D.J. Lydiate, C.P. Smith, J.M. Ward, and H. Schrempf (1985), Genetic Manipulation of Streptomyces, A Laboratory Manual, The John Innes Foundation, Norwich, UK. Protoplasts of S. lividans were prepared by lysozyme treatment, transformed with plasmid DNA, and selected for resistance to thiostrepton, as described in Hopwood, D.A., M.J. Bibb, K.F. Chater, T. Kieser, C.J. Bruton, H.M. Kieser, D.J. Lydiate, C.P. Smith, J.M. Ward, and H. Schrempf (1985), Genetic Manipulation of Streptomyces, A Laboratory Manual, The John Innes Foundation, Norwich, UK. Transformants were screened for proteolytic or amylolytic activity on LB plates containing 30 ug/ml thiostrepton, and either 1% skim milk or 1% corn starch, respectively. E. coli transformants were grown on YT medium containing 50 ug/ml ampicillin.

Materials

Oligonucleotides were synthesized using an Applied Biosystem 380A DNA synthesizer. Columns, phosphoramidites, and reagents used for oligonucleotide synthesis were obtained from Applied Biosystems, Inc. through Technical Marketing Associates. Oligonucleotides were purified by polyacrylamide gel electrophoresis followed by DEAE cellulose chromatography. Enzymes for digesting and modifying DNA were purchased from New England Biolabs and used according to the supplier's recommendations. Radioisotopes [α -³²P]dATP (3000 Ci/mmol) and [γ -³²P]ATP (~3000 Ci/mmol) were from Amersham. Thiostrepton was donated by Squibb.

EXAMPLE 1 - isolation of DNA

Chromosomal DNA was isolated from Streptomyces as described in Chater, K.F., D.A. Hopwood, T. Kieser, and C.J. Thomson (1982), Gene cloning in Streptomyces, Curr. Topics Microbiol. Immunol., 96:69-95, except that sodium dodecyl sarcosinate (final conc. 0.5%) was substituted for sodium dodecyl sulfate. Plasmid DNA of transformed S. lividans was prepared by an alkaline lysis procedure as set out in Hopwood, D.A., M.J. Bibb, K.F. Chater, T. Kieser, C.J. Bruton, H.M. Kieser, D.J. Lydiate, C.P. Smith, J.M. Ward, and H. Schrempf (1985), Genetic Manipulation of Streptomyces, A Laboratory Manual, The John Innes Foundation, Norwich, UK. Plasmid DNA from E. coli was purified by a rapid boiling method (Holmes, D.S., and M. Quigley (1981), A rapid boiling method for the preparation of bacterial plasmids, Anal. Biochem., 114:193-197). DNA fragments and vectors used for all constructions were separated by electrophoresis on low melting point agarose, and purified from the molten agarose by phenol extraction.

EXAMPLE 2 - Construction of Genomic Library

Chromosomal DNA of *S. griseus* ATCC 15395 was digested to completion of *Bam*HI and fractionated by electrophoresis on a 0.8% low melting point agarose gel. DNA fragments ranging in size from 4 to 12 kilobase pairs (kb) were isolated from the agarose gel. The plasmid vectors pUC18 and pUC19 were digested with *Bam*HI, and treated with calf intestinal alkaline phosphatase (Boehringer Mannheim). The *S. griseus* *Bam*HI fragments (0.3 ug) and vectors (0.6 ug) were ligated in a final volume of 20 ul as described in Maniatis, T., E.F. Fritsch, and J. Sambrook (1982), *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY. Approximately 8000 transformants of HB101 were obtained from each ligation reaction.

EXAMPLE 3 - Subcloning of Protease Gene Fragments

A hybrid *Streptomyces-E. coli* vector was constructed by ligating pIJ702, which had been linearized by *Bam*HI, into the *Bam*HI site of pUC8. The unique *Bgl*II site of this vector was used for subcloning *Bam*HI and *Bgl*II fragments to the protease genes. Other fragments were adapted with *Bam*HI linkers to facilitate ligation into the *Bgl*II site. The hybrid vector, with pUC8 inserted at the *Bam*HI site of pIJ702, was incapable of replicating *Streptomyces*. However, the *E. coli* plasmid could be readily removed prior to transforming *S. lividans* by digestion with *Bam*HI followed by recircularization with T4 ligase.

EXAMPLE 4 - Construction for Testing the sprB Signal Peptide

The 0.4 kb *Sau*3AI-*Nco*I fragment containing the aminoglycoside phosphotransferase gene promoter was isolated from pIJ61 and subcloned into the *Bam*HI and *Nco*I sites of a suitable vector. The *Nco*I site containing the initiator ATG was joined to the *Mlu*I site of the *sprB* signal using two 43-mer oligonucleotides, which reconstructed the amino-terminus of the signal peptide. An amylase gene of *S. griseus* was adapted by ligating a 14-mer *Pst*I linker to a *Sma*I site in the third codon. This removed the signal peptide and restored the amino-terminus of the mature amylase. The *Hae*II site of the *sprB* signal was joined to the *Pst*I site of the amylase subclone using two 26-mer oligonucleotides, which reconstructed the carboxy-terminus of the signal peptide.

EXAMPLE 5 - Hybridization

A 20-mer (5'TTCCC(C/G)AACAACGACTACGG3') oligonucleotide was designed from an amino acid sequence (FPNNDYG) which was common to both proteases. For use as a hybridization probe, the oligonucleotide was end-labelled using T4 polynucleotide kinase (New England Biolabs) and [γ -32P]ATP. Digested genomic or plasmid DNA was transferred to a Hybond-N nylon membrane (Amersham) by electroblotting and hybridized in the presence of formamide (50%) as described in Hopwood, D.A., M.J. Bibb, K.F. Chater, T. Kieser, C.J. Bruton, H.M. Kieser, D.J. Lydiate, C.P. Smith, J.M. Ward, and H. Schrempf (1985), *Genetic Manipulation of Streptomyces, A Laboratory Manual*, The John Innes Foundation, Norwich, UK. The filters were hybridized with the labelled oligonucleotide probe at 30°C for 18h, and washed at 47°C. The *S. griseus* genomic library was screened by colony hybridization as described in Wallace, R.B., M.J. Johnson, T. Hirose, T. Miyake, E.H. Kawashima, and K. Itakura (1981), *The use of synthetic oligonucleotides as hybridization probes, II. Hybridization of oligonucleotides of mixed sequence to rabbit globin DNA*, *Nucl. Acids Res.* 9:879-894.

EXAMPLE 6 - DNA Sequencing

The sequences of *sprA* and *sprB* were determined using a combination of the chemical cleavage sequencing method (Maxam, A., and W. Gilbert (1977), *A new method for sequencing DNA*, *Proc. Natl. Acad. Sci. U.S.A.*, 74:560-564) and the dideoxy sequencing method (Sanger, F., S. Nicklen, and A.R. Coulson (1977), *DNA sequencing with chain terminating inhibitors*, *Proc. Natl. Acad. Sci. U.S.A.*, 74:5463-5467). Restriction fragments were end-labeled using either polynucleotide kinase or the large fragment of DNA Polymerase I (Amersham), with the appropriate radiolabeled nucleoside triphosphate. Labeled fragments were either digested with a second restriction endonuclease or strand-separated, followed by electroelution from a polyacrylamide gel. Subclones were prepared in the M13 bacteriophage and the dideoxy sequencing reactions were run using the -20 universal primer (New England Biolabs). In some areas of strong secondary structure, compressions and polymerase failure necessitated the use of

either inosine (Mills, D.R., and F.R. Kramer (1979), Structure independent nucleotide sequence analysis, Proc. Natl. Acad. Sci. U.S.A., 76:2232-2235) or 7-deazaguanosine (Mizusana, S., S. Nishimura, and F. Seela (1986), Improvement of the dideoxy chain termination method of DNA sequencing by use of deoxy-7-deazoguanosine triphosphate in place of dGTP, Nucleic Acids Res., 14:1319-1324) analogs in the dideoxy reactions to clarify the sequence. The sequence were compiled using the software of DNASTARTM - (Daggett, P.E., and F.R. Blattner (1986), Personal access of sequence databases on personal computers, Nucleic Acids Res., 14:611-619).

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. The DNA signal sequence of Fig. 4A.
2. The DNA signal sequence of Fig. 5A.
3. A vector comprising the signal sequence defined in claim 1 or claim 2 and also a sequence encoding a desired protein fused thereto.
4. The vector of claim 3, which is a plasmid or phage.
5. A transformed prokaryotic cell comprising the vector of claim 3 or claim 4, which is capable of expressing said sequences, as a fusion protein.
6. The cell of claim 5, which is of the genus Streptomyces.
7. The cell of claim 6, which is S. lividans or S. griseus.
8. A method for preparing a desired protein, which comprises culturing the cell of any of claims 5 to 7 in a nutrient medium, the fusion protein being produced as an intermediate and the signal sequence directing secretion of the desired protein from the cell.

Claims for the following Contracting State : ES

1. A process for preparing a DNA vector, comprising introducing the signal sequence of Fig. 4A or Fig. 5A and also a sequence encoding a desired protein fused thereto.
2. The process of claim 1, wherein the vector is a plasmid or phage.
3. A process for preparing a transformed prokaryotic cell, comprising transformation with the vector of claim 1 or claim 2, whereby the cell is capable of expressing said sequences, as a fusion protein.
4. The process of claim 3, wherein the cell is of the genus Streptomyces.
5. The process of claim 4, wherein the cell is S. lividans or S. griseus.
6. A method for preparing a desired protein, which comprises culturing the cell of any of claims 3 to 5 in a nutrient medium, the fusion protein being produced as an intermediate and the signal sequence directing secretion of the desired protein from the cell.

Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. DNA-Signalsequenz von Fig. 4A.
2. DNA-Signalsequenz von Fig. 5A.
3. Vektor, der die in Anspruch 1 oder Anspruch 2 definierte Signalsequenz sowie eine damit verknüpfte Sequenz, die für ein gewünschtes Protein kodiert, umfaßt.

4. Vektor nach Anspruch 3, der ein Plasmid oder Phage ist.
5. Transformierte prokaryotische Zelle, welche den Vektor nach Anspruch 3 oder Anspruch 4 umfaßt, die imstande ist, die genannten Sequenzen als Fusionsprotein zu exprimieren.
6. Zelle nach Anspruch 5 der Gattung Streptomyces.
7. Zelle nach Anspruch 6, die S. lividans oder S. griseus ist.
8. Verfahren zur Herstellung eines gewünschten Proteins, welches das Züchten der Zelle nach irgendeinem der Ansprüche 5 bis 7 in einem Nährmedium umfaßt, wobei das Fusionsprotein als Zwischenprodukt erzeugt wird und die Signalsequenz die Sekretion des gewünschten Proteins aus der Zelle steuert.

Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Herstellung eines DNA-Vektors, welches das Einführen der Signalsequenz von Fig. 4A oder Fig. 5A sowie einer damit verknüpften Sequenz, die für ein gewünschtes Protein kodiert, umfaßt.
2. Verfahren nach Anspruch 1, worin ein Vektor ein Plasmid oder Phage ist.
3. Verfahren zur Herstellung einer transformierten prokaryotischen Zelle, umfassend die Transformation mit dem Vektor nach Anspruch 1 oder Anspruch 2, wodurch die Zelle imstande ist, die genannten Sequenzen als Fusionsprotein zu exprimieren.
4. Verfahren nach Anspruch 3, worin die Zelle von der Gattung Streptomyces ist.
5. Verfahren nach Anspruch 4, worin die Zelle S. lividans oder S. griseus ist.
6. Verfahren zur Herstellung eines gewünschten Proteins, welches das Züchten der Zelle nach irgendeinem der Ansprüche 3 bis 5 in einem Nährmedium umfaßt, wobei das Fusionsprotein als Zwischenprodukt erzeugt wird und die Signalsequenz die Sekretion des gewünschten Proteins aus der Zelle steuert.

Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, GR, IT, LI, LU, NL, SE

1. Séquence d'ADN signal de la figure 4A.
2. Séquence d'ADN signal de la figure 5A.
3. Vecteur comportant la séquence signal définie en revendication 1 ou revendication 2, ainsi que, fusionnée à ce vecteur, une séquence codant pour une protéine voulue.
4. Vecteur selon la revendication 3, qui est un plasmide ou un phage.
5. Cellule procaryote transformée comportant le vecteur selon la revendication 3 ou la revendication 4, capable d'exprimer lesdites séquences sous la forme d'une protéine fusionnée.
6. Cellule selon la revendication 5, du genre Streptomyces.
7. Cellule selon la revendication 6, qui est S. lividans ou S. griseus.
8. Procédé de préparation d'une protéine recherchée, qui comporte la mise en culture de la cellule selon l'une quelconque des revendications 5 à 7 dans un milieu nutritif, la protéine fusionnée étant produite en tant qu'intermédiaire, et la séquence signal dirigeant la sécrétion de la protéine recherchée hors de la cellule.

Revendications pour l'Etat contractant suivant : ES

1. Procédé de préparation d'un vecteur d'ADN comportant l'introduction de la séquence signal de la figure 4A ou de la figure 5A, ainsi que, fusionnée à ce vecteur, une séquence codant pour une protéine
5 voulue.
2. Procédé selon la revendication 1, dans lequel le vecteur est un plasmide ou un phage.
3. Procédé de préparation d'une cellule procaryote transformée, comportant la transformation avec le
10 vecteur selon la revendication 1 ou la revendication 2, grâce auquel la cellule est capable d'exprimer lesdites séquences sous la forme d'une protéine fusionnée.
4. Procédé selon la revendication 3, dans lequel la cellule est du genre Streptomyces.
- 15 5. Procédé selon la revendication 4, dans lequel la cellule est S. lividans ou S. griseus.
6. Procédé de préparation d'une protéine recherchée, qui comporte la mise en culture de la cellule selon l'une quelconque des revendications 3 à 5 dans un milieu nutritif, la protéine fusionnée étant produite
20 en tant qu'intermédiaire, et la séquence signal dirigeant la sécrétion de la protéine recherchée hors de la cellule.

25

30

35

40

45

50

55

FIGURE 1

TYPICAL BIOPHARMACEUTICAL WORKUP

Recombinant Protein
Produced in *E. coli*




FIG.2.





FIG.3.



A

GTGACCCCATCTCATTCCGGGCTCGCGGGCGCGAATCCGGCCTTGGGTACGGGACGGTCCCGTCAACGATTC
 CAGCGTGCAACTTGGCAGGTTACGCGCCACTCCCACTGGGTGAGAACCTCGCGEACCAACGGCCCCACCTCACEC
 -116
 M T F K R F S P L S S T S R
 GACCGGGCGTCCCGCCATACCTCGGAGGATCTCGTGACCTTCAAGCGCTTCTCGCCGCTCAGCAGCACGTCAAG
 -100 -80
 Y A R L L A V A S G L V A A A A L A T P S A V A A
 ATATGCACGGCTCTCGCGGTGGCTCCGGCTGGTGGCGCGCGGCGCTGGCCACCCCTCGGCGGTGCGCGC
 -60
 P E A E S K A T V S Q L A D A S S A I L A A D V A
 TCCGAGGGGAGTCCAAGGCCACCGTTTCGACGCTCGCCGACGCCAGCTCCGCCATCTCGCCGCTGATGTGGC
 -40
 G T A W Y T E A S T G K I V L T A D S T V S K A E
 GGGCACCGCTCGTACACGGAGCGAGCACGGCAAGATCGTCTCACCGCCGACAGCACCGGTGCGAAGGCCGA
 -20
 L A K V S N A L A G S K A K L T V K R A E G K F T
 ACTGGCCAAGGTCAACAACCGCTCGCGGGCTCCAAGGCGAACTGACGGTCAAGCGCGCCGAGGSCAAGTTCA
 20
 P L I A G G E A I T T G G S R C S L G F N V S V N
 CCGCGTATCGCGGGCGCGAGGCCATCACCAACCGGTGGCAGCGCTGTTGCTCGGCTTCAACGTGTCGGTCAA
 40
 G V A H A L T A G H C T N I S A S H S I G T R T G
 CGCGCTCGCCACGCGCTCACCGCGGCCACTGCACCAACATCAGCGCCAGCTGGTCCATCGGCACCGCACCGG
 60
 T S F P N N D Y G I I R H S N P A A A D G R V Y L
 AACCACTTCCCGAACAACGACTACGGCATCATCGCCACTCGAACCCGGCGGGCGCCGACGGCCGGGTCTACCT
 80
 Y N G S Y Q D I T T A G N A F V G Q A V Q R S G S
 GTACAACGGCTCTTACCAGGACATCACGACGGCGGGCAACGCCCTTGTGGGGCAGGCCGTCCAGCGCAGCGCAG
 100 120
 T T G L R S G S V T G L N A T V N Y G S S G I V Y
 CACCACCGGGCTGCGCAGCGGCTCGGTACCGGCTCAACGCCACGGTCAACTACGGTTCCAGCGGGATCGTSTA
 140
 G M I Q T N V C A E P G G D S G G S L F A G S T A L
 CGCATGATCCAGACCAACGTCTGTGCCGAGCCGGTGACAGTGGAGGCTCGCTCTTCCGGGCGACGACCGCTCT
 160
 G L T S G G S G N C R T G G T T F Y Q P V T E A L
 GGGTCTCACTCCGGCGGCACTGCGGACCGCGGCGACCGACGTTCTACCAACCGTCAECGAGGCGCT
 181
 S A Y G A T V L * -41.0
 GAGCGCTACGGGCAACGGTCTGTAGCCGGTGCCACCGGGGCTTCGGGCTGACCGCCGACCGGCGCCCGAAG
 -30.8
 CCGCGCGCAGCGCCACCCCGCGGACCGTGTCTCGCGCGGCTCCGCCCTCGCGGTGCCAGCAACCCACCGTC
 CTTTCCCGGTACGGCGCTGCGCTCGACCCGCATCGCGAAGTTGCCGAGAGTGGCGGCTCGACCGGCACTGC
 TGAAGTCTGCGCTCGCGCCACGGTCCGGTTGCGGCCCGCCGGACCGGACCGCGCTGGGGAAAGCCCTCACT
 CAGCCCGGTTGCGCGCGATGAGGTGCGGATACCAAGCGCAAGGAGGCTTCGGGGTGGGACCTGTGTCTCGTGS
 TCGAC

FIG.4.

FIG.4A.




FIG.4B.





FIG.4C.



B

[illegible]

FIG. 5.

FIG.5A.




FIG.5B.




FIG.5C.



1 S G G D A I Y S S T G R C S L G F H Y R S 20
 ATCTCCGGCGGCGACGGATCTACTCCAGCACCAGGAGCTGCTCGCTCGGCTTCAACGTCCGAG
 40
 G S T Y Y F L T A G H C T D G A T T W W A N S A R
 CGGCAGCAGCTACTACTTCTGACCGCCGGCCACTGCACGGACGGCGGACCACTGGTGGGCGAACTCGGCCG
 60
 T T V L G T T S G S S F P N N D Y G I V R Y T N T
 CACCACGGTGCTCGGCACGACCTCCGGGTGAGCTTCCGAACAACGACTACGGCATCGTGGCTACACCAACAC
 80
 T I P K D G T V G G Q D I T S A A N A T V G M A V
 CACCATTCCCAAGGACGGCAGGTCCGGCGCCAGGACATCACCAGCGCCGCCAACGCCACCGTCGGCATGGCGGT
 100
 T R R G S T T G T H S G S V T A L N A T V N Y G G
 CACCGCGCGCGGCTCCACCACCGGCACCCACAGCGGTTCGGTCACCGCACTCAAGGCCACCGTCAACTACGGGG
 120
 G D V V Y G M I R T N V C A E P G D S G G P L Y S
 CGGCAGCTCGTCTACGGCATGATCCGCACCAACGTGTGCGCGAGCCCGCGACTCCGGCGGCCCGCTCTACTC
 140
 G T R A I G L T S G G S G N C S S G G T T F F Q P
 CGGCACCGCGCGATCGGTCTGACCTCCGGCGGACGGCAACTGCTCCTCCGGCGGCACGACCTTCTTCCAGCC
 160
 V T E A L S A Y G V S V Y *
 185
 GGTCAACCGAGGCGCTGAGCGGTACGGCGTCAGCGTGTAC




FIG.6A.

A

```

sprA  MTFKRFSPLSSTSRYPARLLAVASGLVAAAALATPSAVA
      M KR S S R R AV GL A AALA P A A
sprB  MRIKRTSHRSNAARRVRTTAVLAGLAAVAALAVPTANA

```

FIG.6B.

B

```

sprA  APEAESKATVSQLADASSAILAADVAGTAWYTEASTGKI
      A QL AS A L AD AGTAW
sprB  ETPRTFSAN--QLTAASDAVLGADIAGTAWNIDPQSKRL

sprA  VLTADSTVSKAELAKVSNALAGSKAK-LTVKRAEGKFTPL
      V T DSTVSKAE AG A L R GKFT L
sprB  VVTVDSTVSKAEINQIKKS-AGANADALRIERTPGKFTKL

```

FIG.6C.

C

```

sprA  IAGGEAITTGGSRCSLGFHVSVNGVAHALTAGHCTNIS
      I GG AI RCSLGFHV LTAGHCT
sprB  ISGGDAIYSSSTGRCSLGFHVRS GSTYYFLTAGHCTDGA

sprA  ASWS-----IGTRTGTSPFNNDYGIIRHSNPAAA-
      W GT G SFPNNDYGI R N
sprB  TTWWANSARTTVLGTTSGSSFPNNDYGIVRYTNTTIPK

sprA  DGRVYLYNGSYQDITTAGNAFVGQAVQRSGSTTG LRSG
      DG V G QDIT A NA VG AV R GSTTG SG
sprB  DGTV----GG-QDITSAANATVGMATRRGSTTGTHSG

sprA  SVTGLNATVNYGSSGIVYGMITNVCAEPGDSGGSLFA
      SVT LNATVNYG VYGM I TNVCAEPGDSSG L
sprB  SVTALNATVNYGGGDVVYGMIRTNVCAEPGDSSGPLY

sprA  GSTALGLTSGGSGNCRTGGTTFYQPVTEALSAYGATVL
      G A GLTSGGSGNC GGTTF QPVTEALSAYG V
sprB  GTRAIGLTSGGSGNCSSGGTTFQPVTEALSAYGVSVY

```